

Simple Continued Fractions for Some Irrational Numbers, II

J. O. SHALLIT

Department of Mathematics, University of California, Berkeley, California 94720

Communicated by H. Zassenhaus

Received August 27, 1979; revised March 18, 1980

In this paper we prove a theorem allowing us to determine the continued fraction expansion for $\sum_{k=0}^{\infty} u^{-c(k)}$, where $c(k)$ is any sequence of positive integers that grows sufficiently quickly. As an application, we determine the continued fraction expansion for Liouville's famous transcendental number $\sum_{k=0}^{\infty} m^{-(k+1)!}$.

I. INTRODUCTION

In a previous paper [1], the author discussed the continued fractions for the irrational numbers defined by

$$B(u, \infty) = \sum_{k=0}^{\infty} u^{-2^k} \quad (u \geq 2, \text{ an integer}).$$

In this paper, we prove a more general result concerning the continued fractions for the irrational numbers

$$S(u, \infty) = \sum_{k=0}^{\infty} u^{-c(k)},$$

where $c(k)$ is any sequence of positive integers that grows sufficiently quickly. As an application, we determine the continued fraction for Liouville's number.

II. A THEOREM

Let $\{c(k)\}_{k=0}^{\infty}$ be a sequence of positive integers such that $c(v+1) \geq 2c(v)$ for all $v \geq v'$, where v' is a non-negative integer. Let $d(v) = c(v+1) - 2c(v)$. Define $S(u, v)$ as follows:

$$S(u, v) = \sum_{k=0}^v u^{-c(k)}.$$

Then we have the following:

THEOREM. Suppose $v \geq v'$. If $S(u, v) = [a_0, a_1, a_2, \dots, a_n]$ and n is even, then

$$S(u, v+1) = [a_0, a_1, \dots, a_n, u^{d(v)} - 1, 1, a_n - 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1].$$

Proof. This proof uses the ideas behind the proof of Theorem 1 in [1]. Let $c(v+1) > 2c(v)$, and let

$$S(u, v) = [a_0, a_1, \dots, a_n] = p_n/q_n.$$

We find

$$\begin{aligned} [a_0, a_1, \dots, a_n, u^{d(v)} - 1] &= \frac{(u^{d(v)} - 1)p_n + p_{n-1}}{(u^{d(v)} - 1)q_n + q_{n-1}}, \\ [a_0, a_1, \dots, a_n, u^{d(v)} - 1, 1] &= \frac{u^{d(v)}p_n + p_{n-1}}{u^{d(v)}q_n + q_{n-1}}, \\ [a_0, a_1, \dots, a_n, u^{d(v)} - 1, 1, a_n - 1] &= \frac{(a_n - 1)(u^{d(v)}p_n + p_{n-1}) + (u^{d(v)} - 1)p_n + p_{n-1}}{(a_n - 1)(u^{d(v)}q_n + q_{n-1}) + (u^{d(v)} - 1)q_n + q_{n-1}} \\ &= \frac{u^{d(v)}a_n p_n + a_n p_{n-1} - p_n}{u^{d(v)}a_n q_n + a_n q_{n-1} - q_n}, \\ [a_0, a_1, \dots, a_n, u^{d(v)} - 1, 1, a_n - 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1] &= \frac{(u^{d(v)}p_n + p_{n-1})q_{n-2} + (u^{d(v)}a_n p_n + a_n p_{n-1} - p_n)q_{n-1}}{(u^{d(v)}q_n + q_{n-1})q_{n-2} + (u^{d(v)}a_n q_n + a_n q_{n-1} - q_n)q_{n-1}} \\ &= \frac{u^{d(v)}p_n q_n + 1}{u^{d(v)}q_n^2}. \end{aligned}$$

In this last simplification, we use the fact that

$$p_{n-1}q_{n-2} - p_{n-2}q_{n-1} = 1$$

if n is even, and Eqs. (4)–(7) of [1].

Now

$$\begin{aligned} p_n/q_n &= \sum_{k=0}^v u^{-c(k)} \\ &= u^{-c(v)} \left(1 + u \sum_{k=0}^{v-1} u^{c(v)-c(k)-1} \right) \end{aligned}$$

and $(p_n, q_n) = 1$; hence $q_n = u^{c(v)}$. Therefore,

$$\begin{aligned} & [a_0, a_1, \dots, a_n, u^{d(v)} - 1, 1, a_n - 1, a_{n-1}, a_{n-2}, \dots, a_2, a_1] \\ &= \frac{p_n}{q_n} + \frac{1}{u^{d(v)} q_n^2} \\ &= S(u, v) + u^{-c(v+1)} \\ &= S(u, v+1). \end{aligned}$$

It should be noted that zero elements appear in the expansion of $S(u, v+1)$ if $d(v) = 0$. These zero elements may be removed with the use of the formula

$$[a_0, a_1, \dots, a_k, 0, a_{k+1}, \dots] = [a_0, a_1, \dots, a_k + a_{k+1}, a_{k+2}, \dots].$$

Thus, the theorem holds for $c(v) = 2^v$ (as shown in [1]). This proves the theorem for $c(v+1) \geq 2c(v)$.

One hypothesis of the theorem states that n is even. This is not really much of a restriction, since any finite continued fraction can be expressed in two equivalent forms:

If $a_n \geq 2$,

$$[a_0, a_1, \dots, a_n] = [a_0, a_1, \dots, a_n - 1, 1].$$

If $a_n = 1$,

$$[a_0, a_1, \dots, a_{n-1}, 1] = [a_0, a_1, \dots, a_{n-2}, a_{n-1} + 1],$$

for example, see Hardy and Wright [2]. Thus we can always write a continued fraction for $S(u, v)$ such that n is even.

Repeated application of the theorem gives longer and longer prefixes of the continued fraction for $S(u, \infty)$, since it is easily seen that the expansions for $S(u, v)$ and $S(u, \infty)$ coincide up to the penultimate partial denominator in the expansion for $S(u, v)$.

III. AN APPLICATION

Using the theorem, we calculate the continued fraction expansion for the Liouville transcendental number [3]

$$L(m) = \sum_{k=0}^{\infty} m^{-(k+1)!} \quad (m \text{ an integer } \geq 2).$$

Let $c(k) = (k+1)!$. Then $d(k) = k(k+1)!$. Repeated application of the theorem gives

$$\begin{aligned} S(u, 0) &= m^{-1!} = [0, m] = [0, m-1, 1], \\ S(u, 1) &= m^{-1!} + m^{-2!} = [0, m-1, 1, 0, 1, 0, m-1] \\ &= [0, m-1, m+1], \\ S(u, 2) &= m^{-1!} + m^{-2!} + m^{-3!} \\ &= [0, m-1, m+1, m^2-1, 1, m, m-1]. \end{aligned}$$

We may continue in this fashion. Hence

$$L(m) = [0, m-1, m+1, m^2-1, 1, m, m-1, m^{12}-1, 1, m-2, \dots].$$

ACKNOWLEDGMENTS

In a letter dated October 11, 1979, G. Köhler kindly informed the author that he has independently discovered the main theorem, and has generalized it to some extent.

The author would like to thank T. Cusick, G. Köhler, and M. Mendes France for bringing his attention to the work of M. Kmošek, who has also generalized the main theorem in a master's thesis published in Poland in 1979. A. Schinzel reported on Kmošek's work at a conference on Diophantine approximations at Oberwolfach in May, 1979.

The author thanks the referee for several improvements.

REFERENCES

1. J. SHALLIT, Simple continued fractions for some irrational numbers, *J. Number Theory* **11** (1979), 209–217.
2. G. H. HARDY AND E. M. WRIGHT, "An Introduction to the Theory of Numbers," p. 133. Oxford Univ. Press (Clarendon), London, 1971.
3. J. LIOUVILLE, Sur des classes très-étendues de quantités dont la valeur n'est ni algébrique, ni même réductible à des irrationnelles algébriques, *J. Math. Pures Appl.* (1) **16** (1851), 139.